

IN THE UNITED STATES PATENT & TRADEMARK OFFICE

IN RE APPLICATION OF: :

Toru KOYANAGI et al. : EXAMINER: Patricia L. Morris

SERIAL NO.: 10/589,782 :

FILED: August 17, 2006 : GROUP ART UNIT: 1625

FOR: ANTHRANILAMIDES, PROCESS FOR THE PRODUCTION
THEREOF, AND PEST CONTROLLERS CONTAINING THE
SAME

DECLARATION

COMMISSIONER FOR PATENTS

ALEXANDRIA, VIRGINIA 22313

SIR:

Now comes KEIICHI KISHI who deposes and says:

That my name is KEIICHI KISHI;

That my address is 991-5, Minaminakamaru, Minuma-ku,
Saitama-shi, Saitama, Japan,;

That I know well both the English and Japanese languages;

That the attached English language translation is an accurate translation of Japanese Patent Application No. JP2004-133722 filed on April 28, 2004 to the best of my knowledge and belief;

I hereby declare further that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of this application or any patent issuing thereon.

FURTHER DEPONENT SAITH NOT.

August 19, 2009
Date

Keiichi Kishi
KEIICHI KISHI

Date of Application: April 28, 2004

Application Number: Japanese Patent Application
No. 2004-133722

Applicant(s): Ishihara Sangyo Kaisha, Ltd.

【Type of Document】 PETITION FOR PATENT APPLICATION

【Reference No.】 P2004021

【Addressee】 Commissioner, Patent Office

【International Patent Classification】 C07C235/00

【Inventor(s)】

 【Name】 Toru Koyanagi

 【Address or Residence】 Ishihara Sangyo Kaisha, Ltd.
Chuo Kenkyusho
3-1, Nishi-shibukawa 2-chome,
Kusatsu-shi, Shiga-ken

【Inventor(s)】

 【Name】 Masayuki Morita

 【Address or Residence】 Ishihara Sangyo Kaisha, Ltd.
Chuo Kenkyusho
3-1, Nishi-shibukawa 2-chome,
Kusatsu-shi, Shiga-ken

【Inventor(s)】

 【Name】 Kenichi Nakamoto

 【Address or Residence】 Ishihara Sangyo Kaisha, Ltd.
Chuo Kenkyusho
3-1, Nishi-shibukawa 2-chome,
Kusatsu-shi, Shiga-ken

【Inventor(s)】

 【Name】 Akihiro Hisamatsu

 【Address or Residence】 Ishihara Sangyo Kaisha, Ltd.
Chuo Kenkyusho
3-1, Nishi-shibukawa 2-chome,
Kusatsu-shi, Shiga-ken

【Patent Applicant(s)】

 【Identification No.】 000000354

 【Address or Residence】 3-15, Edobori 1-chome, Nishi-ku,
Osaka-shi, Osaka

【Name or Company Name】 Ishihara Sangyo Kaisha, Ltd.

【Representative】 Fujio Tamura

【Identification of Fees】

【Prepayment account No.】 0056823

【Amount of Payment】 16,000 yen

【List of Submitted Items】

【Name of item】	Claim(s)	1
----------------	----------	---

【Name of item】	Specification	1
----------------	---------------	---

【Name of item】	Abstract	1
----------------	----------	---

- 1 -


【TYPE OF DOCUMENT】

CLAIM(S)

【CLAIM 1】

An anthranilamide compound represented by the formula
(I) or its salt:

5 【KA 1】




wherein R^1 is halogen or alkyl; each of R^2 and R^3 which
are independent of each other, is halogen or $-CF_3$; A is
alkyl substituted by Y; Y is haloalkylthio,
10 haloalkylsulfinyl, haloalkylsulfonyl or cycloalkyl which
may be substituted by halogen or alkyl; m is 1 or 2; and
n is 0 or 1.

【CLAIM 2】


A process for producing an anthranilamide compound
15 represented by the formula (I) or its salt:

【KA 2】




wherein R^1 is halogen or alkyl; each of R^2 and R^3 which
are independent of each other, is halogen or $-CF_3$; A is
5 alkyl substituted by Y; Y is haloalkylthio,
haloalkylsulfinyl, haloalkylsulfonyl or cycloalkyl which
may be substituted by halogen or alkyl; m is 1 or 2; and
n is 0 or 1, which comprises (1) reacting a compound
represented by the formula (II):

10 【KA 3】




wherein R^1 , A and m are as defined above, with a compound
represented by the formula (III):

【KA 4】



wherein R^2 , R^3 and n are as defined above, and Z is a chlorine atom, $-OH$ or C_{1-4} alkoxy, or (2) reacting a compound represented by the formula (IV):

【KA 5】



wherein R^1 , R^2 , R^3 , m and n are as defined above, with a compound represented by the formula (V): $A-NH_2$, wherein A is as defined above.

【CLAIM 3】

A pesticide containing the anthranilamide compound or its salt as defined in Claim 1, as an active ingredient.

【CLAIM 4】

An insecticide, miticide or nematocide containing the anthranilamide compound or its salt as defined in Claim 1, as an active ingredient.

【CLAIM 5】

A method for controlling a pest, which comprises applying an effective amount of the anthranilamide compound or its salt as defined in Claim 1.

【TYPE OF DOCUMENT】 DESCRIPTION

【TITLE OF THE INVENTION】

ANTHRANILAMIDE COMPOUNDS, PROCESS FOR THEIR
PRODUCTION AND PESTICIDES CONTAINING THEM

5 【TECHNICAL FIELD】

The present invention relates to a novel
anthranilamide compound which is useful for an active
ingredient of pesticide.

【BACKGROUND ART】

10 WO03/2422 discloses anthranilamide compounds.
However, chemical structures of such compounds are
different from the chemical structure of the
anthranilamide compound represented by the after-
mentioned formula (I).

15 【PATENT DOCUMENT 1】 International Publication
WO03/24222

【DISCLOSURE OF THE INVENTION】


【PROBLEMS TO BE SOLVED BY THE INVENTION】

For many years, many pesticides have been used, but
20 many of them have various problems such that the effects
are inadequate, their use is restricted as pests have
acquired resistance, etc. Accordingly, it is desired to
develop a novel pesticide substantially free from such
problems, for example, a pesticide capable of controlling
25 various pests which create problems in agricultural and
horticultural fields or a pesticide which is capable of
controlling pests parasitic on animals.

[MEANS OF SOLVING THE PROBLEMS]

The present inventors have conducted various studies on anthranilamide compounds in an effort to find a superior pesticide. As a result, they have found that a novel anthranilamide compound or its salt has an extremely high pesticidal effect against pests at a low dose and have accomplished the present invention. Namely, the present invention relates to an anthranilamide compound represented by the formula (I) or its salt:

[KA 1]



wherein R^1 is halogen or alkyl; each of R^2 and R^3 which are independent of each other, is halogen or $-CF_3$; A is alkyl substituted by Y; Y is haloalkylthio, haloalkylsulfinyl, haloalkylsulfonyl or cycloalkyl which may be substituted by halogen or alkyl; m is 1 or 2; and n is 0 or 1; a process for its production; and a pesticide containing it.

[EFFECT OF THE INVENTION]

The pesticide containing, as an active ingredient,

the novel anthranilamide compound represented by the above formula (I), has a very high pesticidal effect against pests at a low dose.

【BEST MODE(S) FOR CARRYING OUT THE INVENTION】

5 The number of substituents Y in A may be 1 or more, and if more, the respective substituents Y may be the same or different. Further, the positions for substitution of the substituents Y may be any positions.

 As the halogen or halogen as the substituent in R¹,
10 R², R³ or Y, an atom of fluorine, chlorine, bromine or iodine may be mentioned. The number of halogens as substituents may be 1 or more, and if more, the respective halogens may be the same or different. Further, the positions for substitution of such halogens
15 may be any positions.

 The alkyl or alkyl moiety in R¹, A or Y may be linear or branched. As its specific example, C₁₋₆ alkyl such as methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, pentyl or hexyl may be mentioned.

20 The number of alkyl as substituents may be 1 or more, and if more, the respective alkyl may be the same or different. Further, the positions for substitution of such alkyls may be any position.

 As a specific example of cycloalkyl or cycloalkyl
25 moiety in Y, C₃₋₆ cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl may be mentioned.


 The salt of the anthranilamide compound of the above

formula (I) includes all kinds so long as they are agriculturally acceptable. For example, an alkali metal salt such as a sodium salt or a potassium salt; an alkaline earth metal salt such as a magnesium salt or a calcium salt; an ammonium salt such as a dimethylammonium salt or a triethylammonium salt; an inorganic acid salt such as a hydrochloride, a perchlorate, a sulfate or a nitrate; or an organic acid salt such as an acetate or a methanesulfonate, may be mentioned.

10 The anthranilamide compound of the formula (I) may have an isomers such as an optical isomer. In the present invention, each isomer and mixtures of isomers are both included. Further, in the present invention, various isomers other than those mentioned above, may be included within the scope of the common knowledge in this technical field. Further, depending upon the type of such an isomer, the chemical structure may be different from the above-mentioned formula (I), but it is obvious to one skilled in the art that such a structure is in isomeric relation and thus falls within the scope of the present invention.

25 The anthranilamide compound of the above formula (I) or its salt (hereinafter referred to simply of the compound of the present invention) can be produced by the following reactions (A) and (B) and in accordance with a usual method for producing a salt.

【KA 2】



R^1 , R^2 , R^3 , A , m and n are as defined above, and Z is a chlorine atom, $-OH$ or C_{1-4} alkoxy.

5 In a case where Z is a chlorine atom, the reaction (A) can be carried out usually in the presence of a base.

As the base, one or more types may suitably be selected for use from, for example, an alkali metal hydroxide such as sodium hydroxide or potassium hydroxide; an alkali metal carbonate such as sodium carbonate or potassium carbonate; an alkali metal hydride such as sodium hydride or potassium hydride; and a tertiary amine such as trimethylamine, triethylamine,

10

triisopropylamine, diisopropylethylamine, pyridine, 4-dimethylaminopyridine, 2,6-dimethylpyridine, 4-pyrrolidinopyridine, N-methylmorpholine, N,N-dimethylaniline, N,N-diethylaniline, N-ethyl-N-methylaniline, 1,8-diazabicyclo[5.4.0]-7-undecene or 1,4-diazabicyclo[2.2.2]octane. The base may be used in an amount of from 1 to 5 times by mol, preferably from 1 to 2.5 times by mol, to the compound of the formula (II).

In a case where Z is a chlorine atom, the reaction (A) can be carried out in the presence of a solvent, as the case requires. The solvent may be any solvent so long as it is a solvent inert to the reaction. For example, one or more types may suitably be selected for use from, for example, an ether such as diethyl ether, butyl ethyl ether, tetrahydrofuran, dioxane or dimethoxyethane; a halogenated hydrocarbon such as chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane, trichloroethane or dichloroethylene; an aromatic hydrocarbon such as benzene, toluene or xylene; a dipolar aprotic solvent such as acetonitrile, propionitrile, N,N-dimethylformamide, dimethylsulfoxide, hexamethylphosphoric triamide, sulfolane, dimethylacetamide or N-methylpyrrolidone; an ester such as methyl acetate, ethyl acetate or propyl acetate; and a ketone such as acetone, diethyl ketone, methyl ethyl ketone or methyl isobutyl ketone.

In a case where Z is a chlorine atom, the reaction (A) can be carried out usually from -20 to +60°C, preferably from 0 to 30°C. The reaction time is usually from about 1 to 24 hours, preferably from about 2 to 12
5 hours.

In a case where Z is -OH, the reaction (A) can be carried out usually in the presence of a dehydration condensing agent and a solvent.

The dehydration condensing agent may, for example,
10 be a carbodiimide such as N,N'-dicyclohexylcarbodiimide, 1,3-diisopropylcarbodiimide or 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride; or others such as 1,1'-carbonyl-bis-1H-imidazole, phenyl
dichlorophosphate, diethyl phosphorocyanidate, 1,3,5-
15 triaza-2,4,6-triphosphorin-2,2,4,4,6,6-hexachloride, cyanuric chloride, isobutyl chloroformate, chlorosulfonyl isocyanate, or trifluoroacetic anhydride.

The solvent may be any solvent so long as it is inert to the reaction. For example, one or more types
20 may suitably be selected for use from, for example, an ether such as diethyl ether, butyl ethyl ether, tetrahydrofuran, dioxane or dimethoxyethane; a halogenated hydrocarbon such as chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon
25 tetrachloride, dichloroethane, trichloroethane or dichloroethylene; an aromatic hydrocarbon such as benzene, toluene or xylene; a dipolar aprotic solvent

such as acetonitrile, propionitrile, N,N-dimethylformamide, dimethylsulfoxide, hexamethylphosphoric triamide, sulfolane, dimethylacetamide or N-methylpyrrolidone; an ester such as methyl acetate, ethyl acetate or propyl acetate; a ketone such as acetone, diethyl ketone, methyl ethyl ketone or methyl isobutyl ketone; and an aliphatic hydrocarbon such as pentane, hexane, heptane, octane or cyclohexane.

In a case where Z is -OH, the reaction (A) can be carried out usually from -20 to +60°C, preferably from 0 to 30°C. The reaction time is usually from about 0.5 to 24 hours, preferably from about 1 to 12 hours.

In a case where Z is C₁₋₄ alkoxy, the reaction (A) can be carried out usually in the presence of a base and a solvent. As the base, one or more types may suitably be selected for use from, for example, an alkali metal hydride such as sodium hydride or potassium hydride; an alkali metal alkoxide such as sodium methoxide, sodium ethoxide or potassium tertiary butoxide; and a tertiary amine such as trimethylamine, triethylamine, triisopropylamine, diisopropylethylamine, pyridine, 4-dimethylaminopyridine, 2,6-dimethylpyridine, 4-pyrrolidinopyridine, N-methyl morpholine, N,N-dimethylaniline, N,N-diethylaniline, N-ethyl-N-methylaniline, 1,8-diazabicyclo[5.4.0]-7-undecene or 1,4-diazabicyclo[2.2.2]octane. The base may be used in an amount of from 1 to 5 times by mol, preferably from 1 to


2.5 times by mol, to the compound of the formula (II).

The solvent may be any solvent so long as it is inert to the reaction. For example, one or more types may suitably be selected for use from, for example, an
5 ether such as diethyl ether, butyl ethyl ether, tetrahydrofuran, dioxane or dimethoxyethane; an aromatic hydrocarbon such as benzene, toluene or xylene; a dipolar aprotic solvent such as acetonitrile, propionitrile, N,N-dimethylformamide, dimethylsulfoxide, hexamethylphosphoric
10 triamide, sulfolane, dimethylacetamide or N-methylpyrrolidone and an alcohol such as methanol, ethanol, propanol, n-butanol or tert-butanol.

In a case where Z is C₁₋₄ alkoxy, the reaction (A) can be carried out usually from 0 to 120°C, preferably
15 from 20 to 80°C. The reaction time is usually from about 0.5 to 24 hours, preferably from about 1 to 12 hours.

The compound of the above formula (II) or (III) may be a known compound or may be produced in accordance with known reference materials. For example, the compound of
20 the formula (II) can be produced by or in accordance with the method disclosed in Synthesis, 1980, p.505. The compound of the formula (III) can be produced by or in accordance with the method disclosed in schemes 9 to 22 in WO03/24222.

[KA 3]



R¹, R², R³, A, m and n are as defined above.

The reaction (B) can be carried out usually in the
5 presence of a solvent.


The solvent may be any solvent so long as it is
inert to the reaction. For example, one or more types
may suitably be selected for use from, for example, an
ether such as diethyl ether, butyl ethyl ether,
10 tetrahydrofuran, dioxane or dimethoxyethane; a
halogenated hydrocarbon such as chlorobenzene,
dichlorobenzene, dichloromethane, chloroform, carbon
tetrachloride, dichloroethane, trichloroethane or
dichloroethylene; an aromatic hydrocarbon such as

benzene, toluene or xylene; and a dipolar aprotic solvent such as acetonitrile, propionitrile, N,N-dimethylformamide, dimethylsulfoxide, hexamethylphosphoric triamide, sulfolane, dimethylacetamide or N-methylpyrrolidone.

The reaction (B) can be carried out usually from 0 to 120°C, preferably from 20 to 80°C. The reaction time is usually about from 0.5 to 24 hours, preferably from about 1 to 12 hours.

The compound of the above formula (IV) may be a known compound or may be produced in accordance with a known reference material. For example, the compound of the formula (IV) can be produced by or in accordance with the method disclosed in Org. Prep. Proceed. Int., 1993, vol.25, p.585 or the method disclosed in schemes 8 to 10 in WO03/24222.

The compound of the above formula (V) includes a novel compound. Such a compound can be produced by the Gabriel method, and can be produced, for example, in accordance with the following reaction (C).



A is as defined above, and T is halogen, -OSO₂G (G is

a sulfonate residue) or -OH. When T is halogen or -OSO₂G, M is sodium or potassium, and when T is -OH, M is a hydrogen atom. The above sulfonate residue may, for example, be a C₁₋₆ alkyl such as methyl or ethyl; or
5 phenyl which may be substituted by C₁₋₆ alkyl.

In a case where T is halogen or -OSO₂G and M is sodium or potassium, the first step of the reaction (C) can be carried out usually in the presence of a solvent.

The solvent may be any solvent so long as it is
10 inert to the reaction. For example, one or more types may suitably be selected for use from, for example, an ether such as diethyl ether, butyl ethyl ether, tetrahydrofuran, dioxane or dimethoxyethane; an aromatic hydrocarbon such as benzene, toluene or xylene; a dipolar
15 aprotic solvent such as acetonitrile, propionitrile, N,N-dimethylformamide, dimethylsulfoxide, hexamethylphosphoric triamide, sulfolane, dimethylacetamide or N-methylpyrrolidone; and an alcohol such as methanol, ethanol, propanol, n-butanol or tert-butanol.

20 In a case where T is halogen or -OSO₂G and M is sodium or potassium, the first step of the reaction (C) can be carried out usually from 0 to 150°C, preferably from 30 to 110°C. The reaction time is usually from about 0.5 to 24 hours, preferably from about 1 to 12
25 hours.

In a case where T is -OH and M is a hydrogen atom, the first step of the reaction (C) can be carried out

usually by Mitsunobu Method. For example, it can be carried out by using a dialkyl azo dicarboxylate and triphenylphosphine in the presence of a solvent. Each of such a dialkyl azo dicarboxylate and triphenylphosphine
5 may be used usually in an amount approximately equimolar to the compound of the formula (VI). The above dialkyl azo dicarboxylate may, for example, be diethyl azo dicarboxylate or diisopropyl azo dicarboxylate.

The solvent may be any solvent so long as it is
10 inert to the reaction. For example, one or more types may suitably be selected for use from, for example, an ether such as diethyl ether, butyl ethyl ether, tetrahydrofuran, dioxane or dimethoxyethane; a halogenated hydrocarbon such as chlorobenzene,
15 dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane, trichloroethane or dichloroethylene; and an aromatic hydrocarbon such as benzene, toluene or xylene.

In a case where T is -OH and M is a hydrogen atom,
20 the first step of the reaction (C) can be carried out usually from 0 to 80°C, preferably from 20 to 60°C. The reaction time is usually from about 0.5 to 24 hours, preferably from about 1 to 16 hours.

The second step of the reaction (C) can be carried
25 out usually by decomposing the compound of the formula (VIII) by means of hydrazine in the presence of a solvent. Such hydrazine may be used in an amount

approximately equimolar to the compound of the formula (VIII).

The solvent may be any solvent so long as it is inert to the reaction. For example, one or more types may suitably selected for use from, for example, an ether such as diethyl ether, butyl ethyl ether, tetrahydrofuran, dioxane or dimethoxyethane; an aromatic hydrocarbon such as benzene, toluene or xylene; and an alcohol such as methanol, ethanol, propanol, n-butanol or tert-butanol.

The second step of the reaction (C) can be carried out usually from 0 to 140°C, preferably from 30 to 100°C. The reaction time is usually from about 0.5 to 24 hours, preferably from about 2 to 12 hours.

Preferred embodiments of pesticides containing the compounds of the present invention will be described below. The pesticides containing the compounds of the present invention are particularly useful, for example, as agents for controlling various pests which become problematic in the agricultural and horticultural fields, i.e. agricultural and horticultural pesticides, or as agents for controlling pests which are parasitic on animals i.e. pesticides against parasites on animals.

The agricultural and horticultural pesticides containing the compounds of the present invention are useful as an insecticide, a miticide, a nematocide and a soil pesticide, and they are effective for controlling

plant parasitic mites such as two-spotted spider mite (Tetranychus urticae), carmine spider mite (Tetranychus cinnabarinus), kanzawa spider mite (Tetranychus kanzawai), citrus red mite (Panonychus citri), European red mite 5 (Panonychus ulmi), broad mite (Polyphagotarsonemus latus), pink citrus rust mite (Aculops pelekassi) and bulb mite (Rhizoglyphus echinopus); agricultural insect pests such as diamondback moth (Plutella xylostella), cabbage armyworm (Mamestra brassicae), common cutworm (Spodoptera 10 litura), codling moth (Laspeyresia pomonella), bollworm (Heliothis zea), tobacco budworm (Heliothis virescens), gypsy moth (Lymantria dispar), rice leafroller (Cnaphalocrocis medinalis), Adoxophyes sp., summer fruit tortrix (Adoxophyes orana fasciata), peach fruit moth 15 (Carposina niponensis), oriental fruit moth (Grapholita molesta), black cutworm (Agrotis ipsilon), cutworm (Agrotis segetum), colorado potato beetle (Leptinotarsa decemlineata), cucurbit leaf beetle (Aulacophora femoralis), boll weevil (Anthonomus grandis), aphids, 20 planthoppers, leafhoppers, scales, bugs, whiteflies, thrips, grasshoppers, anthomyiid flies, scarabs and ants; plant parasitic nematodes such as root-knot nematodes, cyst nematodes, root-lesion nematodes, rice white-tip nematode (Aphelenchoides besseyi), strawberry bud 25 nematode (Nothotylenchus acris), pine wood nematode (Bursaphelenchus lignicolus); gastropods such as slugs and snails; soil pests such as isopods such as pillbugs

(Armadilidium vulgare) and pillbugs (Porcellio scaber);
hygienic insect pests such as tropical rat mite
(Ornithonyssus bacoti), housefly (Musca domestica), house
mosquito (Culex pipiens) and cockroaches; stored grain
5 insect pests such as angoumois grain moth (Sitotroga
cerealella), adzuki bean weevil (Callosobruchus
chinensis), red flour beetle (Tribolium castaneum) and
mealworms; clothes, house and household insect pests such
as casemaking clothes moth (Tinea pellionella), black
10 carpet beetle (Anthrenus scrophularidae) and subterranean
termites; domestic mites such as mold mite (Tyrophagus
putrescentiae), Dermatophagoides farinae and
Chelacaropsis moorei. Among them, the agricultural and
horticultural pesticides containing the compounds of the
15 present invention are particularly effective for
controlling plant parasitic mites, agricultural insect
pests, plant parasitic nematodes or the like. Further,
they are effective against insect pests having acquired
resistance to organophosphorus, carbamate and/or
20 synthetic pyrethroid insecticides. Moreover, the
compounds of the present invention have excellent
systemic properties, and by the application of the
compounds of the present invention to soil treatment, not
only noxious insects, noxious mites, noxious nematodes,
25 noxious gastropods and noxious isopods in soil but also
foliage pests can be controlled.

Another preferred embodiments of the pesticides

containing compounds of the present invention may be agricultural and horticultural pesticides which collectively control the above-mentioned plant parasitic mites, agricultural insect pests, plant parasitic
5 nematodes, gastropods and soil pests.

The agricultural and horticultural pesticide containing the compound of the present invention, is usually formulated by mixing the compound with various agricultural adjuvants and used in the form of a
10 formulation such as a dust, granules, water-dispersible granules, a wettable powder, a water-based suspension concentrate, an oil-based suspension concentrate, water soluble granules, an emulsifiable concentrate, a soluble concentrate, a paste, an aerosol or an ultra low-volume
15 formulation. However, so long as it is suitable for the purpose of the present invention, it may be formulated into any type of formulation which is commonly used in this field. Such agricultural adjuvants include solid carriers such as diatomaceous earth, slaked lime, calcium
20 carbonate, talc, white carbon, kaoline, bentonite, a mixture of kaolinite and sericite, clay, sodium carbonate, sodium bicarbonate, mirabilite, zeolite and starch; solvents such as water, toluene, xylene, solvent naphtha, dioxane, acetone, isophorone, methyl isobutyl ketone,
25 chlorobenzene, cyclohexane, dimethylsulfoxide, N,N-dimethylformamide, dimethylacetamide, N-methyl-2-pyrrolidone, and alcohol; anionic surfactants and

spreaders such as a salt of fatty acid, a benzoate, an alkylsulfosuccinate, a dialkylsulfosuccinate, a polycarboxylate, a salt of alkylsulfuric acid ester, an alkyl sulfate, an alkylaryl sulfate, an alkyl diglycol ether sulfate, a salt of alcohol sulfuric acid ester, an alkyl sulfonate, an alkylaryl sulfonate, an aryl sulfonate, a lignin sulfonate, an alkyldiphenyl ether disulfonate, a polystyrene sulfonate, a salt of alkylphosphoric acid ester, an alkylaryl phosphate, a styrylaryl phosphate, a salt of polyoxyethylene alkyl ether sulfuric acid ester, a polyoxyethylene alkylaryl ether sulfate, a salt of polyoxyethylene alkylaryl ether sulfuric acid ester, a polyoxyethylene alkyl ether phosphate, a salt of polyoxyethylene alkylaryl phosphoric acid ester, and a salt of a condensate of naphthalene sulfonate with formalin; nonionic surfactants and spreaders such as a sorbitan fatty acid ester, a glycerin fatty acid ester, a fatty acid polyglyceride, a fatty acid alcohol polyglycol ether, acetylene glycol, acetylene alcohol, an oxyalkylene block polymer, a polyoxyethylene alkyl ether, a polyoxyethylene alkylaryl ether, a polyoxyethylene styrylaryl ether, a polyoxyethylene glycol alkyl ether, a polyethylene glycol, a polyoxyethylene fatty acid ester, a polyoxyethylene sorbitan fatty acid ester, a polyoxyethylene glycerin fatty acid ester, a polyoxyethylene hydrogenated castor oil, and a polyoxypropylene fatty acid ester; and

vegetable and mineral oils such as olive oil, kapok oil, castor oil, palm oil, camellia oil, coconut oil, sesame oil, corn oil, rice bran oil, peanut oil, cottonseed oil, soybean oil, rapeseed oil, linseed oil, tung oil, and
5 liquid paraffins. Each of the components as such adjuvants may be one or more suitably selected for use, so long as the purpose of the present invention can thereby be accomplished. Further, various additives which are commonly used, such as a filler, a thickener,
10 an anti-settling agent, an anti-freezing agent, a dispersion stabilizer, a phytotoxicity reducing agent, and an anti-mold agent, may also be employed.

The weight ratio of the compound of the present invention to the various agricultural adjuvants is
15 usually from 0.001:99.999 to 95:5, preferably from 0.005:99.995 to 90:10.

In the actual application of such a formulation, it may be used as it is, or may be diluted to a predetermined concentration with a diluent such as water,
20 and various spreaders e.g. surfactants, vegetable oils or mineral oils may be added thereto, as the case requires.

The application of the agricultural and horticultural pesticide containing the compound of the present invention can not generally be defined, as it varies
25 depending upon the weather conditions, the type of the formulation, the application season, the application site or the types or degree of outbreak of the pest insects.

However, it is usually applied in a concentration of the active ingredient being from 0.05 to 800,000 ppm, preferably from 0.5 to 500,000 ppm, and the dose per unit area is such that the compound of the present invention
5 is from 0.05 to 50,000 g, preferably from 1 to 30,000 g, per hectare. Further, agricultural and horticultural pesticides as another preferred embodiment of pesticides containing the compounds of the present invention may be applied in accordance with the above-described
10 application of pesticides. The present invention includes such a method for controlling pests, particularly for controlling plant parasitic mites, agricultural insect pests or plant parasitic nematodes by such applications.

15 Various formulations of agricultural and horticultural pesticides containing the compounds of the present invention or their diluted compositions may be applied by conventional methods for application which are commonly employed, such as spraying (e.g. spraying,
20 jetting, misting, atomizing, powder or grain scattering or dispersing in water), soil application (e.g. mixing or drenching), surface application (e.g. coating, powdering or covering) or impregnation to obtain poisonous feed. Further, it is possible to feed domestic animals with a
25 food containing the above active ingredient and to control the outbreak or growth of pests, particularly insect pests, with their excrements. Furthermore, the

active ingredient may also be applied by a so-called ultra low-volume application method. In this method, the composition may be composed of 100% of the active ingredient.

5 Further, the agricultural and horticultural pesticides containing compounds of the present invention may be mixed with or may be used in combination with other agricultural chemicals, fertilizers or phytotoxicity-reducing agents, whereby synergistic
10 effects or activities may sometimes be obtained. Such other agricultural chemicals include, for example, a herbicide, an insecticide, a miticide, a nematocide, a soil pesticide, a fungicide, an antiviral agent, an attractant, an antibiotic, a plant hormone and a plant
15 growth regulating agent. Especially, with a mixed pesticide having a compound of the present invention mixed with or used in combination with one or more active compounds of other agricultural chemicals, the application range, the application time, the pesticidal
20 activities, etc. may be improved to preferred directions. The compound of the present invention and the active compounds of other agricultural chemicals may separately be formulated so that they may be mixed for use at the time of application, or they may be formulated together.
25 The present invention includes such a mixed pesticidal composition.

The mixing ratio of the compound of the present

invention to the active compounds of other agricultural chemicals can not generally be defined, since it varies depending upon the weather conditions, the types of formulations, the application time, the application site, 5 the types or degree of outbreak of insect pests, etc., but it is usually within a range of from 1:300 to 300:1, preferably from 1:100 to 100:1, by weight. Further, the dose for the application is such that the total amount of the active compounds is from 0.1 to 50,000 g, preferably 10 from 1 to 30,000 g, per hectare. The present invention includes a method for controlling pests by an application of such a mixed pesticide composition.

The active compounds of insect pest control agents such as insecticides, miticides, nematocides or soil 15 pesticides in the above-mentioned other agricultural chemicals, include, for example, (by common names, some of them are still in an application stage) organic phosphate compounds such as Profenofos, Dichlorvos, Fenamiphos, Fenitrothion, EPN, Diazinon, Chlorpyrifos- 20 methyl, Acephate, Prothiofos, Fosthiazate, Phosphocarb, Cadusafos, Disulfoton, Chlorpyrifos, Demeton-S-methyl, Dimethoate and Methamidophos; carbamate compounds such as Carbaryl, Propoxur, Aldicarb, Carbofuran, Thiodicarb, Methomyl, Oxamyl, Ethiofencarb, Pirimicarb, Fenobucarb, 25 Carbosulfan, and Benfuracarb; nereistoxin derivatives such as Cartap, and Thiocyclam, and Bensultap; organic chlorine compounds such as Dicofol and Tetradifon;

organometallic compounds such as Fenbutatin Oxide;
pyrethroid compounds such as Fenvalerate, Permethrin,
Cypermethrin, Deltamethrin, Cyhalothrin, Tefluthrin,
Ethofenprox, Fenpropathrin and Bifenthrin; benzoylurea
5 compounds such as Diflubenzuron, Chlorfluazuron,
Teflubenzuron, Flufenoxuron, Lufenuron and Novaluron;
juvenile hormone-like compounds such as Methoprene,
Pyriproxyfen, and Fenoxycarb; pyridazinone compounds such
as Pyridaben; pyrazole compounds such as Fenpyroximate,
10 Fipronil, Tebufenpyrad, Ethiprole, Tolfenpyrad and
Acetoprole; neonicotinoids such as Imidacloprid,
Nitenpyram, Acetamiprid, Thiacloprid, Thiamethoxam,
Clothianidin, and Dinotefuran; hydrazine compounds such
as Tebufenozide and Methoxyfenozide, Chromafenozide;
15 dinitro compounds; organic sulfur compounds; urea
compounds; triazine compounds; hydrazone compounds; and
other compounds, such as Flonicamid, Buprofezin,
Hexythiazox, Amitraz, Chlordimeform, Silafluofen,
Triazamate, Pymetrozine, Pyrimidifen, Chlorfenapyr,
20 Indoxacarb, Acequinocyl, Etoxazole, Cyromazine, 1,3-
dichloropropene, Diafenthiuron, Benclothiaz, Flufenerim,
Pyridalyl, Spirodiclofen, Bifenazate, Spiromesifen,
Propargite, Clofentezine, Etoxazole and Fluacrypyrim.
Further, microbial agricultural chemicals such as BT
25 agents, insect viruses, entomopathogenic fungi, and
nematophagous fungi, antibiotics such as Avermectin,
Emamectin-Benzate, Milbemectin, Spinosad and Ivermectin,

may be used in admixture or in combination.

The active compounds of fungicides among the above-mentioned other agricultural chemicals include, for example, (by common names, some of which are still in an application stage) pyrimidinamine compounds such as Mepanipyrim, Pyrimethanil, and Cyprodinil; azole compounds such as Triadimefon, Bitertanol, Triflumizole, Etaconazole, Propiconazole, Penconazole, Flusilazole, Myclobutanil, Cyproconazole, Terbuconazole, Hexaconazole, Furconazole-cis, Prochloraz, Metconazole, Epoxiconazole, Tetraconazole, Oxpoconazole, and Sipconazole; quinoxaline compounds such as Quinomethionate; dithiocarbamate compounds such as Maneb, Zineb, Mancozeb, Polycarbamate, Propineb; organic chlorine compounds such as Fthalide, Chlorothalonil, and Quintozene; imidazole compounds such as Benomyl, Thiophanate-Methyl, Carbendazim, and Cyazofamid; pyridinamine compounds such as Fluazinam; cyanoacetamide compounds such as Cymoxanil; phenylamide compounds such as Metalaxyl, Oxadixyl, Ofurace, Benalaxyl, Furalaxyl, and Cyprofuram; sulfenic acid compounds such as Dichlofluanid; copper compounds such as cupric hydroxide, and Oxine Copper; isoxazole compounds such as Hydroxyisoxazole; organophosphorus compounds such as Fosetyl-Al, Tolclofos-Methyl, S-benzyl O,O-diisopropylphosphorothioate, O-ethyl S,S-diphenylphosphorodithioate, and aluminummethylhydrogen phosphonate; N-halogenothioalkyl compounds such as Captan,

Captafol, and Folpet; dicarboximide compounds such as Procymidone, Iprodione, and Vinclozolin; benzanilide compounds such as Flutolanil, Mepronil, and Zoxamide; piperazine compounds such as Triforine; pyridine

5 compounds such as Pyrifenox; carbinol compounds such as Fenarimol; and Flutriafol; piperidine compounds such as Fenpropidine; morpholine compounds such as Fenpropimorph; organotin compounds such as Fentin Hydroxide, and Fentin Acetate; urea compounds such as Pencycuron; cinnamic acid

10 compounds such as Dimethomorph; phenylcarbamate compounds such as Diethofencarb; cyanopyrrole compounds such as Fludioxonil, and Fenpiclonil; Strobilurin compounds such as Azoxystrobin, Kresoxim-Methyl, Metominofen, Trifloxystrobin, Picoxystrobin, and Pyraclostrobin;

15 oxazolidinedione compounds such as Famoxadone; thiazole carboxamide compounds such as Ethaboxam; silyl amide compounds such as Silthiopham; aminoacid amidecarbamate compounds such as Iprovalicarb; imidazolidine compound such as Fenamidone; hydroxyanilide compounds such as

20 Fenhexamid; benzene sulfonamide compounds such as Flusulfamide; anthraquinone compounds; crotonic acid compounds; antibiotics; and other compounds, such as Isoprothiolane, Tricyclazole, Pyroquilon, Diclomezine, Probenazole, Quinoxifen, Propamocarb Hydrochloride,

25 Spiroxamine, Chloropicrin, Dazomet, and Metam-Sodium.

Further, agricultural chemicals which may be used in admixture with or in combination with the compounds of

the present invention, may, for example, be the active ingredient compounds in the herbicides as disclosed in Farm Chemicals Handbook (2002 edition), particularly those of soil treatment type.

5 The pesticides against parasites on animals are effective for controlling e.g. external parasites which are parasitic on the body surface of host animals (such as the back, the axilla, the lower abdomen or inside of the thigh) or internal parasites which are parasitic in
10 the body of host animals (such as the stomach, the intestinal tract, the lung, the heart, the liver, the blood vessels, the subcutis or lymphatic tissues), but they are particularly effective for controlling the external parasites.

15 The external parasites may, for example, be animal parasitic acarina or fleas. Their species are so many that it is difficult to list all of them, and therefore, their typical examples will be given.

 The animal parasitic acarina may, for example, be
20 ticks such as Boophilus microplus, Rhipicephalus sanguineus, Haemaphysalis longicornis, Haemaphysalis flava, Haemaphysalis campanulata, Haemaphysalis concinna, Haemaphysalis japonica, Haemaphysalis kitaokai, Haemaphysalis ias, Ixodes ovatus, Ixodes nipponensis,
25 Ixodes persulcatus, Amblyomma testudinarium, Haemaphysalis megaspinosa, Dermacentor reticulatus, and Dermacentor taiwanesis; common red mite (Dermanyssus

gallinae); northern fowl mites such as Ornithonyssus
sylviarum, and Ornithonyssus bursa; trombidoids such as
Eutrombicula wichmanni, Leptotrombidium akamushi,
Leptotrombidium pallidum, Leptotrombidium fuji,
5 Leptotrombidium tosa, Neotrombicula autumnalis,
Eutrombicula alfreddugesi, and Helenicula miyagawai;
cheyletidae such as Cheyletiella yasguri, Cheyletiella
parasitivorax, and Cheyletiella blakei; sarcoptic mange
mites such as Psoroptes cuniculi, Chorioptes bovis,
10 Otodectes cynotis, Sarcoptes scabiei, and Notoedres cati;
and Demodicidae such as Demodex canis. The pesticides
against parasites on animals, containing the compounds of
the present invention, are particularly effective for the
control of ticks among them.

15 The fleas may, for example, be externally parasitic
wingless insects belonging to Siphonaptera, more
specifically, fleas belonging to Pulicidae, Ceratephyllus,
etc. Fleas belonging to Pulicidae may for example, be
Ctenocephalides canis, Ctenocephalides felis, Pulex
20 irritans, Echidnophaga gallinacea, Xenopsylla cheopis,
Leptopsylla segnis, Nosopsyllus fasciatus, and
Monopsyllus anisus. The pesticides against parasites on
animals, containing the compounds of the present
invention, are particularly effective for the control of
25 fleas belonging to Pulicidae, particularly
Ctenocephalides canis and Ctenocephalides felis, among
them.

Other external parasites may, for example, be sucking lice (Anoplura) such as shortnosed cattle louse (Haematopinus eurysternus), horse sucking louse (Haematopinus asini), sheep louse, longnosed cattle louse 5 (Linognathus vituli), and head louse (Pediculus capitis); biting lice such as dog biting louse (Trichodectes canis); and blood-sucking dipterous insects such as horsefly (Tabanus trigonus), biting midges (Culicoides schultzei), and blackfly (Simulium ornatum). Further, 10 the internal parasites may, for example, be nematodes such as lung worms, taenia, trematoda, Fasciola hepatica; and protozoa such as coccidia, malaria parasites (Plasmodium malariae), intestinal sarcocyst, toxoplasma, and cryptosporidium.

15 The host animals may, for example, be pet animals such as dogs, cats, mice, rats, hamsters, guinea pigs, squirrels, rabbits, ferrets, birds (such as pigeons, parrots, hill mynas, Java sparrows, honey parrots, lovebirds and canaries); domestic animals such as cows, 20 horses, pigs and sheep; and poultry such as ducks and chickens. The pesticides against parasites on animals, containing the compounds of the present invention, are particularly effective for the control of pests parasitic on pet animals or domestic animals, especially for the 25 control of external parasites, among them. Among pet animals or domestic animals, they are effective particularly for dogs, cats, cows and horses.

When the compound of the present invention is used as a pesticide against parasites on animals, it may be used as it is or may be used together with suitable adjuvants, as formulated into various formulations such as a dust, 5 granules, tablets, a powder, capsules, a soluble concentrate, an emulsifiable concentrate, a water-based suspension concentrate and an oil-based suspension concentrate. In addition to such formulations, it may be formulated into any type of formulation which is commonly 10 used in this field, so long as it is suitable for the purpose of the present invention. The adjuvants to be used for formulations may, for example, be anionic surfactants or nonionic surfactants exemplified above as adjuvants for formulation of agricultural and 15 horticultural pesticides; a cationic surfactant such as cetyl trimethylammonium bromide; a solvent such as water, acetone, acetonitrile, monomethylacetamide, dimethylacetamide, dimethylformamide, 2-pyrrolidone, N-methyl-2-pyrrolidone, kerosene, triacetin, methanol, 20 ethanol, isopropanol, benzyl alcohol, ethylene glycol, propylene glycol, polyethylene glycol, liquid polyoxyethylene glycol, butyl diglycol, ethylene glycol monomethyl ether, ethylene glycol monoethyl ether, diethylene glycol monoethyl ether, diethylene glycol n- 25 butyl ether, dipropylene glycol monomethyl ether, or dipropylene glycol n-butyl ether; an antioxidant such as butylhydroxyanisole, butylhydroxytoluene, ascorbic acid,

sodium hydrogenmetasulfite, propyl gallate or sodium thiosulfate; a coating film-forming agent such as polyvinylpyrrolidone, polyvinyl alcohol, or a copolymer of vinyl acetate and vinyl pyrrolidone; the vegetable
5 oils and mineral oils as exemplified above as adjuvants for formulation of agricultural and horticultural pesticides; and a carrier such as lactose, sucrose, glucose, starch, wheat flour, corn powder, soybean cake and meal, defatted rice bran, calcium carbonate or other
10 commercially available feed materials. One or more of the respective components of these adjuvants may be suitably selected for use, so long as such will not depart from the purpose of the present invention. Further, other than the above-mentioned adjuvants, some
15 among those known in this field may suitably be selected for use, and still further, some among the above-mentioned various adjuvants to be used in the agricultural and horticultural field may suitably be selected for use.

20 The blend ratio of the compound of the present invention to various adjuvants is usually from 0.1:99.9 to 90:10. In the actual use of such a formulation, it may be used as it is, or may be diluted to a predetermined concentration with a diluent such as water,
25 and various spreaders (e.g. surfactants, vegetable oils or mineral oils) may be added thereto, as the case requires.

Administration of the compound of the present invention to a host animal is carried out orally or parenterally. As an oral administration method, a method of administering a tablet, a liquid agent, a capsule, a
5 wafer, a biscuit, a minced meat or other feed, containing the compound of the present invention, may be mentioned. As a parenteral administration method, there may, for example, be mentioned a method wherein the compound of the present invention is formulated into a suitable
10 formulation and then taken into the body by e.g. intravenous administration, intramuscular administration, intradermal administration, hypodermic administration, etc.; a method wherein it is administered on the body surface by spot-on treatment, pour-on treatment or spray
15 treatment; or a method of embedding a resin fragment or the like containing the compound of the present invention under the skin of the host animal.

The dose of the compound of the present invention to a host animal varies depending upon the administration
20 method, the purpose of administration, the diseased symptom, etc., but it is usually administered in a proportion of from 0.01 mg to 100 g, preferably from 0.1 mg to 10 g, per 1 kg of the body weight of the host animal.

25 The present invention also includes a method for controlling a pest by the above-mentioned administration method or by the above-mentioned dose, particularly a

method for controlling external parasites or internal parasites.

When the compound of the present invention is used as a pesticide against parasites on animals, various
5 vitamins, minerals, amino acids, nutrients, enzymes, antipyretics, sedatives, antiphlogistics, fungicides, colorants, aromatic substances, preservatives, etc., may be used in admixture with or in combination with the adjuvants. Further, as the case requires, other animal
10 drugs or agricultural chemicals, such as vermicides, anti-coccidium agents, insecticides, miticides, pulicides, nematocides, bactericides or antibacterial agents, may be mixed or combined for use, whereby improved effects may sometimes be obtained. The present invention includes
15 such a mixed pesticidal composition having the above-mentioned various components mixed or combined for use, and further a method for controlling a pest by using it, particularly a method for controlling external parasites or internal parasites.

20 **【EXAMPLE(S)】**

Now, the present invention will be described with reference to Examples, but it should be understood that the present invention is by no means limited thereto. Firstly, Preparation Examples of the compound of the
25 present invention will be described.

PREPARATION EXAMPLE 1

Preparation of N-[6-[[cyclopropylmethyl]amino]carbonyl]-

2-methylphenyl]-1-(3-chloro-2-pyridyl)-3-(trifluoromethyl)-1H-pyrazol-5-carboxamide (after-mentioned compound No. 2)

1.49 g of triethylamine was gradually dropwise added
5 to a mixed solution comprising 0.8 g of cyclopropylmethanamine hydrochloride and 40 ml of tetrahydrofuran under cooling with ice, followed by stirring at room temperature for 30 minutes. Then, a mixed solution comprising 1 g of 2-[1-(3-chloro-2-
10 pyridyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl]-8-methyl-4H-3,1-benzoxazin-4-one and 10 ml of tetrahydrofuran was gradually dropwise added. After completion of the dropwise addition, the mixed solution was reacted for 4 hours under reflux. After completion of the reaction,
15 the solvent was distilled off under reduced pressure, and to the residue, ethyl acetate and water were added for extraction. The organic layer was washed with water and a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. The solvent was
20 distilled off under reduced pressure, and the residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate=1/1) to obtain 0.54 g of the desired product having a melting point of 199.4°C.

PREPARATION EXAMPLE 2

25 Preparation of N-[4-chloro-6-[(2-difluoromethylthioethyl)amino]carbonyl]-2-methylphenyl]-1-(3-chloro-2-pyridyl)-3-(trifluoromethyl)-1H-pyrazol-5-

carboxamide (after-mentioned compound No. 3)

(1) 8 g of difluoromethylthio ethyl acetate was gradually dropwise added to a 100 ml of diethyl ether suspension containing 1.08 g of lithium aluminum hydride at room temperature. Then, the mixed solution was reacted for 4 hours under reflux. After completion of the reaction, the reaction mixture was cooled with ice, and 2.16 ml of water and 1.72 ml of a 10% sodium hydroxide aqueous solution were dropwise added thereto, followed by stirring at room temperature for one night. Insolubles were filtered off, and the solvent was distilled off under reduced pressure to obtain 4.2 g of oily 2-(difluoromethylthio)ethanol.

(2) 5.93 g of diisopropyl azo dicarboxylate was gradually dropwise added to a mixed solution comprising 3.76 g of 2-(difluoromethylthio)ethanol obtained in the above step, 4.32 g of phthalimide, 7.69 g of triphenylphosphine and 100 ml of tetrahydrofuran at room temperature, and the mixed solution was reacted for one night at room temperature. After completion of the reaction, the solvent was distilled off under reduced pressure, and the residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate=4/1) to obtain 5.12 g of N-2-(difluoromethylthio)ethylphthalimide. NMR spectrum data of the above compound is as follows.

$^1\text{H-NMR}$ δ ppm (solvent: CDCl_3 /300 MHz)

3.13 (t, 2H), 3.98 (t, 1H), 6.84 (t, 2H), 7.75 (m, 2H),

7.85 (m, 2H)

- (3) 4.98 g of N-2-(difluoromethylthio)ethylphthalimide obtained in the above step and 1.23 g of hydrazine hydrate (80% solution) were dissolved in 50 ml of ethanol, and the mixed solution was reacted for 5 hours under reflux. The reaction solution was cooled, 2 ml of concentrated hydrochloric acid was added thereto, and the mixed solution was further reacted. A formed precipitate was filtered off, and filtrate was concentrated. 100 ml of water was added to the residue, insolubles were filtered off, and water was distilled off under reduced pressure to obtain 3.14 g of 2-(difluoromethylthio)ethylamine hydrochloride having a melting point of 70.8°C.
- (4) 1.37 g of triethylamine was gradually dropwise added to a mixed solution comprising 1.11 g of 2-(difluoromethylthio)ethylamine hydrochloride obtained in the above step and 50 ml of tetrahydrofuran under cooling with ice, followed by stirring at room temperature for 30 minutes. Then, a mixed solution comprising 1 g of 6-chloro-2-[1-(3-chloro-2-pyridyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl]-8-methyl-4H-3,1-benzoxazin-4-one and 10 ml of tetrahydrofuran was gradually dropwise added thereto. After completion of the dropwise addition, the mixed solution was reacted for 5 hours under reflux. After completion of the reaction, the solvent was distilled off under reduced pressure, and to the residue, ethyl acetate

and water were added for extraction. The organic layer was washed with water and a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. The solvent was distilled off under reduced
5 pressure, and the residue was purified by silica gel column chromatography (eluent: n-hexane/ethyl acetate=1/1) to obtain 0.43 g of the desired product having a melting point of 192.0°C.

Now, typical examples of the compound of the present
10 invention represented by the above formula (I) will be given in Table 1. These compounds can be prepared by the above-described Preparation Examples or by the above-mentioned various processes for the production of the compound of the present invention.

15 In Table 1, No. represents compound No. Further, in Table 1, Me represents a methyl group and CPr a cyclopropyl group. Further, in Table 1, 2-Me-CPr represents a cyclopropyl group having a methyl group substituted at the 2-position. The same applies to other
20 similar descriptions.

【TABLE 1】




TABLE 1

No.	R ^{1a}	R ^{1b}	R ^{1c}	R ^{1d}	R ²	R ³	A	Physical properties (m.p.:°C)
1	Me	H	H	H	CF ₃	Cl	CH ₂ CH(Me)SCHF ₂	216.9
2	Me	H	H	H	CF ₃	Cl	CH ₂ -[CPr]	199.4
3	Me	H	Cl	H	CF ₃	Cl	CH ₂ CH ₂ SCHF ₂	192.0
4	Me	H	Cl	H	CF ₃	Cl	CH ₂ -[CPr]	216.4
5	Me	H	Cl	H	CF ₃	Cl	CH ₂ CH(Me)SCHF ₂	188.3
6	Me	H	Cl	H	CF ₃	Cl	CH(Me)-[CPr]	210.0
7	Me	H	Cl	H	CF ₃	Cl	CH ₂ SCHF ₂	
8	Me	H	Cl	H	CF ₃	Cl	CH(Me)SCHF ₂	
9	Me	H	Cl	H	CF ₃	Cl	CH ₂ C(Me) ₂ SCHF ₂	
10	Me	H	Cl	H	CF ₃	Cl	CH ₂ -[2-Me-CPr]	
11	Me	H	Cl	H	CF ₃	Cl	CH ₂ CH ₂ SCF ₃	
12	Me	H	Cl	H	Cl	Cl	CH ₂ CH ₂ SCHF ₂	226.1
13	Me	H	Cl	H	Br	Cl	CH ₂ CH ₂ SCHF ₂	
14	Me	H	Br	H	CF ₃	Cl	CH ₂ CH ₂ SCHF ₂	
15	Br	H	Br	H	CF ₃	Cl	CH ₂ CH ₂ SCHF ₂	
16	Cl	H	Br	H	CF ₃	Cl	CH ₂ CH ₂ SCHF ₂	
17	Br	H	F	H	CF ₃	Cl	CH ₂ CH ₂ SCHF ₂	
18	Br	H	Cl	H	CF ₃	Cl	CH ₂ CH ₂ SCHF ₂	
19	Me	H	Cl	H	CF ₃	Cl	CH ₂ -[2,2-Cl ₂ -1-Me-CPr]	219.3
20	Me	H	Cl	H	Cl	CF ₃	CH ₂ CH ₂ SCHF ₂	
21	Me	H	Cl	H	Cl	Cl	CH ₂ -[CPr]	
22	Me	H	Cl	H	Br	Cl	CH ₂ -[CPr]	
23	Me	H	Cl	H	CF ₃	Cl	CH ₂ CH(Me)SOCHF ₂	
24	Me	H	Cl	H	CF ₃	Cl	CH ₂ CH(Me)SO ₂ CHF ₂	
25	Me	H	Cl	H	CF ₃	Cl	CH ₂ CH ₂ SOCHF ₂	109.7
26	Me	H	Cl	H	CF ₃	Cl	CH ₂ CH ₂ SO ₂ CHF ₂	108.2
27	Me	H	Cl	H	CF ₃	Cl	C(Me) ₂ CH ₂ SOCHF ₂	
28	Me	H	Cl	H	CF ₃	Cl	CH(Me)CH ₂ CH ₂ SOCHF ₂	
29	Me	H	Cl	H	CF ₃	Cl	CH(Me)CH ₂ CH ₂ SO ₂ CHF ₂	
30	Me	H	Cl	H	Cl	Cl	CH(Me)-[CPr]	186.7
31	Me	H	Cl	H	Br	Cl	CH(Me)-[CPr]	183.6
32	Me	H	Cl	H	CF ₃	Cl	CH ₂ -[1-Me-CPr]	161.2
33	Me	H	Br	H	CF ₃	Cl	CH(Me)-[CPr]	163.7
34	Br	H	Br	H	CF ₃	Cl	CH(Me)-[CPr]	165.5
35	Cl	H	Br	H	CF ₃	Cl	CH(Me)-[CPr]	176.5
36	Br	H	Cl	H	CF ₃	Cl	CH(Me)-[CPr]	158.2
37	Me	H	Cl	H	CF ₃	Cl	CH(Me)CH ₂ SCH ₂ CF ₃	149-151
38	Me	H	Cl	H	CF ₃	Cl	CH(Me)CH ₂ SOCH ₂ CF ₃	153-155
39	Me	H	Cl	H	CF ₃	Cl	CH(Me)CH ₂ SO ₂ CH ₂ CF ₃	171-173

Now, test examples will be described.

TEST EXAMPLE 1

Test on controlling effects against common cutworm

(Spodoptera litura)

5 A leaf segment of cabbage was dipped for about 10
seconds in an insecticidal solution prepared to bring the
concentration of the compound of the present invention to
50 ppm and dried in air. A wet filter paper was laid in
a petri dish having a diameter of 9 cm, and the dried
10 leaf segment of cabbage was placed thereon. Ten second-
third instar larvae common cutworm were released therein
and after putting a cover, left in a constant temperature
chamber at 25°C with lightening. Dead larvae were
counted 5 days after the release, and the mortality was
15 calculated by the following equation. The mortality was
obtained with respect to the above-mentioned compound
Nos. 1 to 3, 6, 12, 19, 25, 26 and 30 to 36, whereby all
compounds showed high controlling effects with a
mortality of at least 90%.

20
$$\text{Mortality (\%)} = (\text{number of dead insects} / \text{number of} \\ \text{released insects}) \times 100$$

Now, Formulation Example will be described.

FORMULATION EXAMPLE 1

(1) Compound of the present invention	
25	20 parts by weight
(2) Clay	72 parts by weight
(3) Sodium lignin sulfonate	8 parts by weight

The above components are uniformly mixed to obtain a wettable powder.

FORMULATION EXAMPLE 2

- | | | |
|---|---------------------------------------|--------------------|
| | (1) Compound of the present invention | |
| 5 | | 5 parts by weight |
| | (2) Talc | 95 parts by weight |

The above components are uniformly mixed to obtain a dust.

FORMULATION EXAMPLE 3

- | | | |
|----|---------------------------------------|--------------------|
| 10 | (1) Compound of the present invention | |
| | | 20 parts by weight |
| | (2) N,N'-dimethylacetamide | 20 parts by weight |
| | (3) Polyoxyethylenealkylphenyl ether | |
| | | 10 parts by weight |
| 15 | (4) Xylene | 50 parts by weight |

The above components are uniformly mixed and dissolved to obtain an emulsifiable concentrate.

FORMULATION EXAMPLE 4

- | | | |
|----|--------------------------------------|--------------------|
| | (1) Clay | 68 parts by weight |
| 20 | (2) Sodium lignin sulfonate | 2 parts by weight |
| | (3) Polyoxyethylenealkylaryl sulfate | |
| | | 5 parts by weight |
| | (4) Fine silica powder | 25 parts by weight |

A mixture of the above components is mixed with
25 compound of the present invention in a weight ratio of
4:1 to obtain a wettable powder.

FORMULATION EXAMPLE 5

(1) Compound of the present invention

50 parts by weight

(2) Oxylated polyalkylphenyl

5 phosphate-triethanolamine 2 parts by weight

(3) Silicone 0.2 part by weight

(4) Water 47.8 parts by weight

The above components are uniformly mixed and
pulverized to obtain a base liquid, and

10 (5) Sodium polycarboxylate 5 parts by weight

(6) Anhydrous sodium sulfate 42.8 parts by weight

are added, and the mixture is uniformly mixed, granulated
and dried to obtain water-dispersible granules.

FORMULATION EXAMPLE 6

15 (1) Compound of the present invention

5 parts by weight

(2) Polyoxyethyleneoctylphenyl ether

1 part by weight

(3) polyoxyethylene phosphoric

20 acid ester 0.1 part by weight

(4) Granular calcium carbonate 93.9 parts by weight

The above components (1) to (3) are preliminarily
uniformly mixed and diluted with a proper amount of
acetone, and then the mixture is sprayed onto the
25 component (4), and acetone is removed to obtain granules.

FORMULATION EXAMPLE 7

(1) Compound of the present invention

2.5 parts by weight

(2) N-methyl-2-pyrrolidone 2.5 parts by weight

(3) Soybean oil 95.0 parts by weight

The above components are uniformly mixed and
5 dissolved to obtain an ultra low volume formulation.

FORMULATION EXAMPLE 8

(1) Compound of the present invention

40 parts by weight

(2) Oxylated polyalkylphenylphosphate-

10 triethanolamine 2 parts by weight

(3) Silicone 0.2 part by weight

(4) Zanthan gum 0.1 part by weight

(5) Ethylene glycol 5 parts by weight

(6) Water 52.7 parts by weight

15 The above components are uniformly mixed and
pulverized to obtain a water-based suspension concentrate.

FORMULATION EXAMPLE 9

(1) Compound of the present invention

10 parts by weight

20 (2) Diethylene glycol monoethyl

ether 90 parts by weight

The above components are uniformly mixed to obtain a
soluble concentrate.

【TYPE OF DOCUMENT】

ABSTRACT

【SUMMARY】


【OBJECT】

For many years, many pesticides have been used, but
5 many of them have various problems such that the effects
are inadequate, their use is restricted as pests have
acquired resistance, etc. Accordingly, it is desired to
develop a novel pesticide substantially free from such
problems.

10 【MEANS OF ACCOMPLISHING THE OBJECT】

The present inventors have various studies on
anthranilamide compounds and as a result, have
accomplished the present invention. Namely, the present
invention is an anthranilamide compound represented by
15 the formula (I) or its salt:

【KA 1】



wherein R¹ is halogen or alkyl; each of R² and R³ which
are independent of each other, is halogen or -CF₃; A is
20 alkyl substituted by Y; Y is haloalkylthio,
haloalkylsulfinyl, haloalkylsulfonyl or cycloalkyl which

may be substituted by halogen or alkyl; m is 1 or 2; and
n is 0 or 1.

【SELECTED FIGURE】

NO SELECTED FIGURE